



Unmasking fungal pathogens by studying MAPK-dependent cell wall regulation in *Candida albicans*

Paul J. Cullen & Mira Edgerton

To cite this article: Paul J. Cullen & Mira Edgerton (2016): Unmasking fungal pathogens by studying MAPK-dependent cell wall regulation in *Candida albicans*, *Virulence*, DOI: [10.1080/21505594.2016.1177695](https://doi.org/10.1080/21505594.2016.1177695)

To link to this article: <http://dx.doi.org/10.1080/21505594.2016.1177695>



Accepted author version posted online: 18 Apr 2016.
Published online: 18 Apr 2016.



[Submit your article to this journal](#)



Article views: 224



[View related articles](#)



[View Crossmark data](#)

EDITORIAL

Unmasking fungal pathogens by studying MAPK-dependent cell wall regulation in *Candida albicans*

Paul J. Cullen^a and Mira Edgerton^b

^aDepartment of Biological Sciences, State University of New York at Buffalo, Buffalo, NY, USA; ^bDepartment of Oral Biology, State University of New York at Buffalo, Buffalo, NY, USA

ARTICLE HISTORY Received 1 April 2016; Revised 6 April 2016; Accepted 8 April 2016

KEYWORDS fungal pathogenesis; glycosylation; immune recognition; MAP kinase; receptor

Fungal infections are a worldwide health problem and are a leading cause of death in developing countries. They contribute to pneumonia,¹ asthma,² and blindness.³ Fungal infections also compound the morbidity of other diseases like tuberculosis.⁴ In developed countries, invasive fungal infections underlie mortality in immune-compromised individuals with HIV/AIDS⁵ and cancer patients undergoing chemotherapy,⁶ and are the fourth largest cause of blood infections. Improved diagnosis and new anti-fungal drugs are recognized as key routes to improving patient survival.⁷ Identifying and increasing the efficacy of anti-fungal drugs is a world-wide health research priority.⁸

The cell wall is an attractive therapeutic target for fungal pathogens. This extracellular organelle is accessible to anti-fungal drugs and is the first site of contact by the immune system. The current picture of the cell wall comes from extensive research in multiple areas, including studies in model organisms where genetic and biochemical approaches have identified cell wall proteins and polysaccharides.⁹ The major polysaccharides in the cell wall are β -1,3-glucans consisting of long linear chains of β -1,3-linked glucose. Less abundant polysaccharides include chitin and β -1,6-glucan. There is extensive cross-linking between the polysaccharides that comprise the cell wall. The cell wall is also composed of glycoproteins, which are glycosylated by covalently modified N- and O-linked oligosaccharide addition.^{10,11} The principle oligosaccharide modification of cell wall proteins is mannosylation. Mannosylation can impact the function, stability, and localization of proteins and can also lead to the cross linking of glycoproteins to polysaccharides in the cell wall.¹²

The cell wall is a dynamic structure. It is rigid enough to provide support and protection, yet is malleable enough to allow for growth and differentiation. What's

more, the cell wall can be reorganized in response to environmental perturbations to promote adaptability and disguise cells from recognition by the immune system.¹³ Among the pathways that regulate cell wall biosynthesis and remodeling are mitogen-activated protein kinase (MAPK) pathways. MAPK pathways are evolutionarily conserved signaling pathways that control stress responses, cell proliferation, and cell differentiation in eukaryotes.^{14,15} In *C. albicans*, one MAPK pathway implicated in cell wall regulation is the Cek1 pathway.¹⁶

The Cek1 pathway is regulated by transmembrane sensors that detect alterations in the environment and initiate cellular responses to these changes. The mucin-like glycoprotein Msb2 is one of the regulators of the Cek1 pathway. Msb2 is a single-pass transmembrane protein with a cytosolic signaling domain and a heavily glycosylated extracellular domain.^{17,18} The extracellular domain of Msb2 is processed by secreted aspartic proteases (SAPs). Release of the inhibitory glycosylated domain is required for MAPK signaling.^{19,20} Msb2 detects (by some mechanism) limiting nutrients in the environment and can also sense elevated temperatures.²¹

Other signaling pathways also impact cell wall integrity. One is the cell wall integrity or Protein Kinase C (PKC) pathway.²²⁻²⁴ Another is the Unfolded Protein Response (UPR) pathway,^{21,25-27} whose major function is to control protein folding in the endoplasmic reticulum.²⁸ Not surprisingly, problems with folding and glycosylation of secreted and cell wall associated proteins can impact the integrity of the cell wall. The fact that multiple pathways regulate the cell wall underscores its critical role in regulating fungal homeostasis.

Landmark discoveries have opened up the field by identifying receptors in the immune system and their substrates

CONTACT Paul J. Cullen ✉ pjcullen@buffalo.edu Department of Biological Sciences, State University of New York at Buffalo, Buffalo, NY 14260-1300, USA.; Mira Edgerton ✉ edgerto@buffalo.edu Department of Oral Biology, State University of New York at Buffalo, Buffalo, NY 14260-1300, USA.

Comment on: Román E, et al. The Cek1 mediated MAP kinase pathway regulates exposure of alpha 1,2 and β 1,2 mannosides in the cell wall of *Candida albicans* modulating immune recognition. Virulence 2016; <http://dx.doi.org/10.1080/21505594.2016.1163458>

© 2016 Taylor & Francis

on the fungal surface.²⁹ Mannose-binding lectin is a serum protein of the innate immune system that binds to and promotes clearance of fungal microorganisms.³⁰ The dectin-1 receptor on macrophages recognizes β -1,3-glucan in the fungal cell wall.^{31,32} The dectin-2 receptor is a pattern recognition receptor on dendritic and other cells that recognizes fungal α -mannan.^{33,34} The pattern recognition receptor galectin-3 on macrophages detects phospholipomannan,³⁵ a sphingolipid in *C. albicans*³⁶ that when recognized elicits a pathogen-specific response.³⁷ Importantly, dectin-1 and galectin-3 interact synergistically to improve the outcome of host immune response to *C. albicans* β -glucans.³⁸

In this issue, Román et al.³⁹ build on the picture of Cek1 pathway-mediated control of cell wall regulation in *C. albicans*. They show that cells lacking an intact Cek1 pathway (*cek1* mutant) exhibit increased exposure of α -1,2-mannosides and β -1,2-mannosides. They reach this conclusion by examining cell wall proteins with antibodies against α -1,2- and β -1,2-mannosides. Under different conditions and treatments, the authors see a striking increase in the electrophoretic mobility of proteins with these modifications between wild-type cells and the *cek1* mutant. This may indicate that cell wall proteins become under-glycosylated in the *cek1* mutant or alternatively fail to become linked to the cell wall. Together with the fact that the *cek1* mutant has abnormally appearing cell walls by transmission electron microscopy, and that *cek1* mutants have increased β -1,3-glucan exposure,^{40,41} their findings support the idea that the cell wall is generally disorganized in cells lacking the Cek1 pathway. This idea is supported by a second set of experiments. The authors show that tunicamycin, an inhibitor of N-linked glycosylation that triggers the UPR pathway and leads to under-glycosylation of cell wall glycoproteins,⁴² exacerbates the growth and cell wall defects of the *cek1* mutant. Their conclusion is in line with other reports^{18,21,43} and may be due to the misfolding, misincorporation, and/or problems with the processing and release of under-glycosylated cell wall proteins.⁴⁴

The authors go on to test the role of the immune system in Cek1-mediated cell wall control. Because galectin-3 can bind β -1,2-mannosides,³⁷ the authors examined whether Gal3^{-/-} mice showed increased fungal susceptibility in a Cek1-dependent manner. As expected from previous studies,¹⁶ *C. albicans* lacking the Cek1 pathway showed reduced virulence in systemically infected mice. However, mice lacking Gal3 did not restore the virulence of the *cek1* mutant. Unexpectedly, Gal3^{-/-} mice survived even better than wild-type mice systemically infected with *C. albicans*. Thus, this work from the Pla group points to gaps in understanding the relationship between Gal3 and the molecular basis of *C. albicans* recognition in the immune response.

This study by Román et al.³⁹ identifies directions for future research. An important question moving forward is to understand how the Cek1 pathway produces a cell wall that is 'invisible' to the immune system. The CEK pathway helps maintain an organized and healthy cell wall by regulating proteins that control cell-wall biosynthesis and remodeling. Who are these proteins? One class of proteins are O-mannosyltransferases (Pmt proteins) that initiate O-mannosylation and selectively glycosylate mucins and other cell wall proteins.^{41,45,46} Other target proteins continue to be uncovered by gene expression profiling by the Pla group and other laboratories.⁴⁷ Biochemical analysis of these cell wall remodeling enzymes may elucidate the precise role the CEK pathway plays in regulating the cell wall. Innovative approaches, like multivariate analysis of morphometric data,⁴⁸ may accelerate progress in this area. These proteins themselves represent attractive drug targets, because they function outside the cell and may be accessible to anti-fungal compounds.

A second question has to do with understanding the functions of cell wall regulatory pathways. At least 3 major pathways control the cell wall: Cek1, PKC, and UPR. Genetic buffering between these pathways complicates assignment of the functions of each pathway in regulating cell wall function. Further complicating matters is the fact that Cek1, PKC, and UPR show cross-regulation.^{21,49-51} Therefore, new approaches are needed to decipher the precise functions of the pathways in regulating the cell wall. Ultimately, studies on MAPK-dependent cell wall regulation and immune recognition promises to elucidate the functions of MAPK pathways and their unique targets with the long-term goals of reducing virulence and improving human health.

Disclosure of potential conflicts of interest

No potential conflicts of interest were disclosed.

Funding

Apologies to authors whose work was not cited due to page limitations. The authors are funded by grant from NIDCR (DE022720).

References

- [1] Pirrone M, Pinciroli R, Berra L. Microbiome, biofilms, and pneumonia in the ICU. *Curr Opin Infect Dis* 2016; 29:160-6; PMID:26855081; <http://dx.doi.org/10.1097/QCO.0000000000000255>
- [2] Ogawa H, Fujimura M, Ohkura N, Satoh K, Makimura K. Fungus-associated asthma: overcoming challenges in diagnosis and treatment. *Expert Rev Clin Immunol* 2014;

- 10:647-56; PMID:24580109; <http://dx.doi.org/10.1586/1744666X.2014.892829>
- [3] Kredics L, Narendran V, Shobana CS, Vagvolgyi C, Manikandan P, Indo-Hungarian Fungal Keratitis Working G. Filamentous fungal infections of the cornea: a global overview of epidemiology and drug sensitivity. *Mycoses* 2015; 58:243-60; PMID:25728367; <http://dx.doi.org/10.1111/myc.12306>
- [4] Lawn SD, Bekker LG, Middelkoop K, Myer L, Wood R. Impact of HIV infection on the epidemiology of tuberculosis in a peri-urban community in South Africa: the need for age-specific interventions. *Clin Infect Dis* 2006; 42:1040-7; PMID:16511773; <http://dx.doi.org/10.1086/501018>
- [5] Armstrong-James D, Meintjes G, Brown GD. A neglected epidemic: fungal infections in HIV/AIDS. *Trends Microbiol* 2014; 22:120-7; PMID:24530175; <http://dx.doi.org/10.1016/j.tim.2014.01.001>
- [6] d'Enfert C. Hidden killers: persistence of opportunistic fungal pathogens in the human host. *Curr Opin Microbiol* 2009; 12:358-64; PMID:19541532; <http://dx.doi.org/10.1016/j.mib.2009.05.008>
- [7] Hahn-Ast C, Glasmacher A, Muckter S, Schmitz A, Kraemer A, Marklein G, Brossart P, von Lilienfeld-Toal M. Overall survival and fungal infection-related mortality in patients with invasive fungal infection and neutropenia after myelosuppressive chemotherapy in a tertiary care centre from 1995 to 2006. *J Antimicrob Chemother* 2010; 65:761-8; PMID:20106864; <http://dx.doi.org/10.1093/jac/dkp507>
- [8] Krysan DJ. Toward improved anti-cryptococcal drugs: Novel molecules and repurposed drugs. *Fungal Genet Biol* 2015; 78:93-8; PMID:25514636; <http://dx.doi.org/10.1016/j.fgb.2014.12.001>
- [9] Free SJ. Fungal cell wall organization and biosynthesis. *Adv Genet* 2013; 81:33-82; PMID:23419716; <http://dx.doi.org/10.1016/B978-0-12-407677-8.00002-6>
- [10] Klis FM, Sosinska GJ, de Groot PW, Brul S. Covalently linked cell wall proteins of *Candida albicans* and their role in fitness and virulence. *FEMS Yeast Res* 2009; 9:1013-28; PMID:19624749; <http://dx.doi.org/10.1111/j.1567-1364.2009.00541.x>
- [11] Chaffin WL. *Candida albicans* cell wall proteins. *Microbiol Mol Biol Rev* 2008; 72:495-544; PMID:18772287; <http://dx.doi.org/10.1128/MMBR.00032-07>
- [12] Ao J, Chinnici JL, Maddi A, Free SJ. The N-Linked Outer Chain Mannans and the Dfg5p and Dcw1p Endo-alpha-1,6-Mannanases Are Needed for Incorporation of *Candida albicans* Glycoproteins into the Cell Wall. *Eukaryot Cell* 2015; 14:792-803; PMID:26048011; <http://dx.doi.org/10.1128/EC.00032-15>
- [13] Halme A, Bumgarner S, Styles C, Fink GR. Genetic and epigenetic regulation of the FLO gene family generates cell-surface variation in yeast. *Cell* 2004; 116:405-15; PMID:15016375; [http://dx.doi.org/10.1016/S0092-8674\(04\)00118-7](http://dx.doi.org/10.1016/S0092-8674(04)00118-7)
- [14] Dhillon AS, Hagan S, Rath O, Kolch W. MAP kinase signalling pathways in cancer. *Oncogene* 2007; 26:3279-90; PMID:17496922; <http://dx.doi.org/10.1038/sj.onc.1210421>
- [15] Qi M, Elion EA. MAP kinase pathways. *J Cell Sci* 2005; 118:3569-72; PMID:16105880; <http://dx.doi.org/10.1242/jcs.02470>
- [16] Csank C, Schroppel K, Leberer E, Harcus D, Mohamed O, Meloche S, Thomas DY, Whiteway M. Roles of the *Candida albicans* mitogen-activated protein kinase homolog, Cek1p, in hyphal development and systemic candidiasis. *Infect Immun* 1998; 66:2713-21; PMID:9596738
- [17] Cullen PJ, Sabbagh W, Jr., Graham E, Irick MM, van Olden EK, Neal C, Delrow J, Bardwell L, Sprague GF, Jr. A signaling mucin at the head of the Cdc42- and MAPK-dependent filamentous growth pathway in yeast. *Genes Dev* 2004; 18:1695-708; PMID:15256499; <http://dx.doi.org/10.1101/gad.1178604>
- [18] Roman E, Cottier F, Ernst JF, Pla J. Msb2 signaling mucin controls activation of Cek1 mitogen-activated protein kinase in *Candida albicans*. *Eukaryot Cell* 2009; 8:1235-49; PMID:19542310; <http://dx.doi.org/10.1128/EC.00081-09>
- [19] Vadaie N, Dionne H, Akajagbor DS, Nickerson SR, Krysan DJ, Cullen PJ. Cleavage of the signaling mucin Msb2 by the aspartyl protease Yps1 is required for MAPK activation in yeast. *J Cell Biol* 2008; 181:1073-81; PMID:18591427; <http://dx.doi.org/10.1083/jcb.200704079>
- [20] Puri S, Kumar R, Chadha S, Tati S, Conti HR, Hube B, Cullen PJ, Edgerton M. Secreted aspartic protease cleavage of *Candida albicans* Msb2 activates Cek1 MAPK signaling affecting biofilm formation and oropharyngeal candidiasis. *PLoS One* 2012; 7:e46020; PMID:23139737; <http://dx.doi.org/10.1371/journal.pone.0046020>
- [21] Saraswat D, Kumar R, Pande T, Edgerton M, Cullen PJ. Signaling Mucin Msb2 Regulates Adaptation to Thermal Stress in *Candida albicans*. *Mol Microbiol* 2016; PMID:26749104; 100(3):425-41
- [22] Levin DE. Regulation of cell wall biogenesis in *Saccharomyces cerevisiae*: the cell wall integrity signaling pathway. *Genetics* 2011; 189:1145-75; PMID:22174182; <http://dx.doi.org/10.1534/genetics.111.128264>
- [23] Navarro-Garcia F, Alonso-Monge R, Rico H, Pla J, Sentandreu R, Nombela C. A role for the MAP kinase gene MKC1 in cell wall construction and morphological transitions in *Candida albicans*. *Microbiology* 1998; 144 (Pt 2):411-24; PMID:9493378; <http://dx.doi.org/10.1099/00221287-144-2-411>
- [24] Navarro-Garcia F, Sanchez M, Pla J, Nombela C. Functional characterization of the MKC1 gene of *Candida albicans*, which encodes a mitogen-activated protein kinase homolog related to cell integrity. *Mol Cell Biol* 1995; 15:2197-206; PMID:7891715; <http://dx.doi.org/10.1128/MCB.15.4.2197>
- [25] Adhikari H, Cullen PJ. Metabolic Respiration Induces AMPK- and Ire1p-Dependent Activation of the p38-Type HOG MAPK Pathway. *PLoS Genet* 2014; 10:e1004734; PMID:25356552; <http://dx.doi.org/10.1371/journal.pgen.1004734>
- [26] Leach MD, Cowen LE. Surviving the heat of the moment: a fungal pathogens perspective. *PLoS Pathog* 2013; 9:e1003163; PMID:23505364; <http://dx.doi.org/10.1371/journal.ppat.1003163>
- [27] Wimalasena TT, Enjalbert B, Guillemette T, Plumridge A, Budge S, Yin Z, Brown AJ, Archer DB. Impact of the unfolded protein response upon genome-wide expression patterns, and the role of Hac1 in the polarized growth, of *Candida albicans*. *Fungal Genet Biol* 2008; 45:1235-47; PMID:18602013; <http://dx.doi.org/10.1016/j.fgb.2008.06.001>

- [28] Cox JS, Shamu CE, Walter P. Transcriptional induction of genes encoding endoplasmic reticulum resident proteins requires a transmembrane protein kinase. *Cell* 1993; 73:1197-1206; PMID:8513503; [http://dx.doi.org/10.1016/0092-8674\(93\)90648-A](http://dx.doi.org/10.1016/0092-8674(93)90648-A)
- [29] Netea MG, Brown GD, Kullberg BJ, Gow NA. An integrated model of the recognition of *Candida albicans* by the innate immune system. *Nat Rev Microbiol* 2008; 6:67-78; PMID:18079743; <http://dx.doi.org/10.1038/nrmicro1815>
- [30] Brouwer N, Dolman KM, van Houdt M, Sta M, Roos D, Kuijpers TW. Mannose-binding lectin (MBL) facilitates opsonophagocytosis of yeasts but not of bacteria despite MBL binding. *J Immunol* 2008; 180:4124-32; PMID:18322223; <http://dx.doi.org/10.4049/jimmunol.180.6.4124>
- [31] Brown GD, Gordon S. Immune recognition. A new receptor for beta-glucans. *Nature* 2001; 413:36-7; PMID:11544516; <http://dx.doi.org/10.1038/35092620>
- [32] Brown GD. Dectin-1: a signalling non-TLR pattern-recognition receptor. *Nat Rev Immunol* 2006; 6:33-43; PMID:16341139; <http://dx.doi.org/10.1038/nri1745>
- [33] Hall RA, Gow NA. Mannosylation in *Candida albicans*: role in cell wall function and immune recognition. *Mol Microbiol* 2013; 90:1147-61; PMID:24125554; <http://dx.doi.org/10.1111/mmi.12426>
- [34] Saijo S, Ikeda S, Yamabe K, Kakuta S, Ishigame H, Akitsu A, Fujikado N, Kusaka T, Kubo S, Chung SH, et al. Dectin-2 recognition of alpha-mannans and induction of Th17 cell differentiation is essential for host defense against *Candida albicans*. *Immunity* 2010; 32:681-91; PMID:20493731; <http://dx.doi.org/10.1016/j.immuni.2010.05.001>
- [35] Jouault T, Iбата-Ombetta S, Takeuchi O, Trinel PA, Sacchetti P, Lefebvre P, Akira S, Poulain D. *Candida albicans* phospholipomannan is sensed through toll-like receptors. *J Infect Dis* 2003; 188:165-72; PMID:12825186; <http://dx.doi.org/10.1086/375784>
- [36] Fradin C, Bernardes ES, Jouault T. *Candida albicans* phospholipomannan: a sweet spot for controlling host response/inflammation. *Semin Immunopathol* 2015; 37:123-30; PMID:25394861; <http://dx.doi.org/10.1007/s00281-014-0461-5>
- [37] Kohatsu L, Hsu DK, Jegalian AG, Liu FT, Baum LG. Galectin-3 induces death of *Candida* species expressing specific beta-1,2-linked mannans. *J Immunol* 2006; 177:4718-26; PMID:16982911; <http://dx.doi.org/10.4049/jimmunol.177.7.4718>
- [38] Esteban A, Popp MW, Vyas VK, Strijbis K, Ploegh HL, Fink GR. Fungal recognition is mediated by the association of dectin-1 and galectin-3 in macrophages. *Proc Natl Acad Sci U S A* 2011; 108:14270-5; PMID:21825168; <http://dx.doi.org/10.1073/pnas.1111415108>
- [39] Román E, Correia I, Salazin A, Fradin C, Jouault T, Poulain D, Liu F-T, Pla J. The Cek1 mediated MAP kinase pathway regulates exposure of alpha 1,2 and beta 1,2 mannosides in the cell wall of *Candida albicans* modulating immune recognition. *Virulence* 2016
- [40] Galan-Diez M, Arana DM, Serrano-Gomez D, Kremer L, Casasnovas JM, Ortega M, Cuesta-Dominguez A, Corbi AL, Pla J, Fernandez-Ruiz E. *Candida albicans* beta-glucan exposure is controlled by the fungal CEK1-mediated mitogen-activated protein kinase pathway that modulates immune responses triggered through dectin-1. *Infect Immun* 2010; 78:1426-36; PMID:20100861; <http://dx.doi.org/10.1128/IAI.00989-09>
- [41] Cantero PD, Ernst JF. Damage to the glycoshield activates PMT-directed O-mannosylation via the Msb2-Cek1 pathway in *Candida albicans*. *Mol Microbiol* 2011; 80:715-25; PMID:21375589; <http://dx.doi.org/10.1111/j.1365-2958.2011.07604.x>
- [42] Sanz P, Herrero E, Sentandreu R. Role of glycosylation in the incorporation of intrinsic mannoproteins into cell walls of *Saccharomyces cerevisiae*. *FEMS Microbiol Lett* 1989; 57:265-269; PMID:2656386; <http://dx.doi.org/10.1111/j.1574-6968.1989.tb03346.x>
- [43] Herrero de Dios C, Roman E, Diez C, Alonso-Monge R, Pla J. The transmembrane protein Opy2 mediates activation of the Cek1 MAP kinase in *Candida albicans*. *Fungal Genet Biol* 2013; 50:21-32; PMID:23149115; <http://dx.doi.org/10.1016/j.fgb.2012.11.001>
- [44] Miller KA, DiDone L, Krysan DJ. Extracellular secretion of overexpressed glycosylphosphatidylinositol-linked cell wall protein Utr2/Chr2p as a novel protein quality control mechanism in *Saccharomyces cerevisiae*. *Eukaryot Cell* 2010; 9:1669-79; PMID:20833895; <http://dx.doi.org/10.1128/EC.00191-10>
- [45] Gentsch M, Tanner W. The PMT gene family: protein O-glycosylation in *Saccharomyces cerevisiae* is vital. *EMBO J* 1996; 15:5752-9; PMID:8918452
- [46] Szafranski-Schneider E, Swidergall M, Cottier F, Tielker D, Roman E, Pla J, Ernst JF. Msb2 shedding protects *Candida albicans* against antimicrobial peptides. *PLoS Pathog* 2012; 8:e1002501; PMID:22319443; <http://dx.doi.org/10.1371/journal.ppat.1002501>
- [47] Huang H, Marcus D, Whiteway M. Transcript profiling of a MAP kinase pathway in *C. albicans*. *Microbiol Res* 2008; 163:380-93; PMID:18467082; <http://dx.doi.org/10.1016/j.micres.2008.03.001>
- [48] Okada H, Ohnuki S, Roncero C, Konopka JB, Ohya Y. Distinct roles of cell wall biogenesis in yeast morphogenesis as revealed by multivariate analysis of high-dimensional morphometric data. *Mol Biol Cell* 2014; 25:222-33; PMID:24258022; <http://dx.doi.org/10.1091/mbc.E13-07-0396>
- [49] Scrimale T, Didone L, de Mesy Bentley KL, Krysan DJ. The unfolded protein response is induced by the cell wall integrity mitogen-activated protein kinase signaling cascade and is required for cell wall integrity in *Saccharomyces cerevisiae*. *Mol Biol Cell* 2009; 20:164-75; PMID:18971375; <http://dx.doi.org/10.1091/mbc.E08-08-0809>
- [50] Adhikari H, Vadaie N, Chow J, Caccamise LM, Chavel CA, Li B, Bowitch A, Stefan CJ, Cullen PJ. Role of the unfolded protein response in regulating the mucin-dependent filamentous-growth mitogen-activated protein kinase pathway. *Mol Cell Biol* 2015; 35:1414-32; PMID:25666509; <http://dx.doi.org/10.1128/MCB.01501-14>
- [51] Roman E, Alonso-Monge R, Miranda A, Pla J. The Mkk2 MAPKK Regulates Cell Wall Biogenesis in Cooperation with the Cek1-Pathway in *Candida albicans*. *PLoS One* 2015; 10:e0133476; PMID:26197240; <http://dx.doi.org/10.1371/journal.pone.0133476>